

Two new neolignans from *Manglietia insignis*

Shan-Zhai SHANG,^{a,b} Ju-Ming YAN,^{a,b} Hai-Bo ZHANG,^a Yi-Ming SHI,^{a,b} Zhong-Hua GAO,^{a,b} Xue DU,^a Yan LI,^a Wei-Lie XIAO,^{a,*} and Han-Dong SUN^{a,*}

^aState Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan, China

^bUniversity of Chinese Academy of Sciences, Beijing 100049, China

Received 26 July 2012; Accepted 17 October 2012

© The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract: Two new neolignans, manneoinsignins A (**1**) and B (**2**), together with four known lignans, were isolated from the leaves and stems of *Manglietia insignis*. The new compounds were established on the basis of extensive spectroscopic analyses. All compounds except **2** were tested for their cytotoxic activity. Compound **3** showed weak cytotoxic activity against the HL-60 human tumor cell line with the IC₅₀ value of 23.5 μM.

Keywords: Magnoliaceae, *Manglietia insignis*, neolignan, cytotoxic activity

Introduction

Plants of the family Magnoliaceae are mainly distributed in southeastern Asia, which contain more than 250 species and are a rich source of lignans possessing various pharmacological functions.^{1–9} Structural and biological diversity of lignans in magnoliaceae family prompted us to investigate *Manglietia insignis* Rehd. et Wils., a plant widely distributed in the west of China and being partly used as a substitute of *M. officinalis* in Yunnan and Sichuan provinces of China.^{4,10} As a result, two new neolignans, manneoinsignins A (**1**) and B (**2**), together with other four known lignans, scaphopetalone (**3**),¹¹ mesosecoisolariciresinol (**4**),¹² lariciresinol (**5**)¹³ and evafolin B (**6**),¹⁴ were isolated. Herein, the isolation, structural elucidation, and their cytotoxic activity are described.

Results and Discussion

Manneoinsignin A (**1**) was obtained as yellow gum. The negative ESIMS of **1** showed characteristic peaks for [M – H][–] and [M – H + 2][–] with the ratio being 3:1, which suggested the presence of a chlorine atom. The molecular formula of **1** was determined as C₁₈H₁₉O₄Cl by its HRESIMS with a pseudomolecular ion peak at *m/z* 333.0897 [M – H][–], corresponding to nine unsaturation degrees. The ¹H NMR spectrum displayed signals due to two sets of ABX-type aromatic systems (δ_H 7.28, 6.91 and 7.24; δ_H 7.08, 6.84 and 7.02), one terminal double bond (δ_H 5.00, 5.05 and 5.97), two methylenes (δ_H 3.33 and 3.61), and two methines (δ_H 4.67 and




Figure 1. The structures of compounds 1–6

3.85) (Table 1). The ¹³C NMR (DEPT) spectrum exhibited eighteen carbon signals, consisting of fourteen *sp*² ones (two of which were signals of allyl group at δ_C 115.6 and 139.4), two methylenes, and two oxygenated methines (Table 1). Comparison of 1D NMR spectra of **1** with those of magnolol showed that they possessed the same skeleton.¹⁵ The differences could be rationalized to be that the allyl group in magnolol was replaced by a 1,2-dihydroxy-3-chlorine-propyl group in **1**. This was further supported by the HMBC correlations of H-7 (δ_H 4.67, 1H, d, *J* = 5.9 Hz) with C-1 (δ_C 134.2), C-2 (δ_C 131.3), C-6 (δ_C 128.2), C-8 (δ_C 76.9) and C-9 (δ_C 47.3), together with the ¹H-¹H COSY correlations of H-7/H-8/H₂-9 (Fig. 2). Furthermore, the HMBC correlations of H-7 with C-1, C-2 and C-6 indicated that the 1,2-dihydroxy-3-chlorine-propyl group was located at C-1 (Fig. 2). The chlorine atom was deduced to be located at C-9 by the characteristic chemical shift of C-9 at δ_C 47.3.


*To whom correspondence should be addressed. E-mail: xwl@mail.kib.ac.cn (W.L. Xiao); hdsun@mail.kib.ac.cn (H.D. Sun)

Table 1. ^1H and ^{13}C NMR spectroscopic assignments of compounds **1** and **2**^a

| position | 1 | | 2 | |
|----------|--|--------------------------------|--|--------------------------------|
| | $\delta_{\text{H}}^{\text{b}}$ | $\delta_{\text{C}}^{\text{c}}$ | $\delta_{\text{H}}^{\text{b}}$ | $\delta_{\text{C}}^{\text{c}}$ |
| 1 | | 134.2 s | | 134.0 s |
| 2 | 7.28 (d, 2.0) | 131.3 d | 7.30 (s) | 131.6 d |
| 3 | | 127.7 s | | 128.4 s |
| 4 | | 155.2 s | | 156.5 s |
| 5 | 6.91 (d, 8.3) | 117.4 d | 6.88 (d, 8.2) | 117.9 d |
| 6 | 7.24 (dd, 8.3, 2.0) | 128.2 d | 7.22 (d, 8.2) | 128.7 d |
| 7 | 4.67 (d, 5.9) | 75.4 d | 4.56 (d, 6.8) | 75.8 d |
| 8 | 3.85 (m) | 76.9 d | 3.88 (m) | 76.5 d |
| 9 | 3.61 (dd, 11.4, 4.0); 3.33 (overlap) | 47.3 t | 3.77 (dd, 11.2, 2.7); 3.67 (dd, 11.2, 6.7) | 48.1 t |
| 1' | | 133.0 s | | 132.6 s |
| 2' | 7.08 (d, 2.0) | 132.7 d | 7.13 (s) | 132.6 d |
| 3' | | 127.6 s | | 128.2 s |
| 4' | | 153.6 s | | 154.7 s |
| 5' | 6.84 (d, 8.2) | 117.4 d | 6.82 (d, 8.1) | 118.0 d |
| 6' | 7.02 (dd, 8.2, 2.0) | 129.8 d | 7.00 (d, 8.1) | 129.8 d |
| 7' | 3.36 (overlap) | 40.4 t | 3.34 (overlap) | 40.7 d |
| 8' | 5.97 (m) | 139.4 d | 5.98 (m) | 139.8 d |
| 9' | 5.05 (dd, 17.0, 1.6); 5.00 (dd, 10.2, 1.6) | 115.6 t | 5.05 (br. d, 17.0); 5.00 (br. d, 9.9) | 115.6 t |

^aRecorded in methanol- d_4 ; ^bRecorded at 600 MHz; ^cRecorded at 125 MHz.

Manneosignin B (**2**) was obtained as yellow gum. The presence of a chlorine atom was deduced by the characteristic peaks for $[\text{M} - \text{H}]^-$ and $[\text{M} - \text{H} + 2]^-$ with the ratio being 3:1 in its negative ESIMS. The molecular formula of **2** was deduced as $\text{C}_{18}\text{H}_{19}\text{O}_4\text{Cl}$ from its HREIMS (m/z 334.0964 $[\text{M}]^+$). Detailed analysis of the 1D NMR data of **1** and **2** suggested that both of them had the same plane structure (Table 1). It was found that the differences of the ^1H NMR chemical shifts of H-7 and H-9 from δ_{H} 4.67, 3.61, and 3.33 in **1** changing to δ_{H} 4.56, 3.77, and 3.67 in **2** and the ^{13}C NMR chemical shifts of C-7, C-8 and C-9 from δ_{C} 75.4, 76.9, and 47.3 in **1** changing to δ_{C} 75.8, 76.5, and 48.1. The reason for the minor distinct NMR data of **1** and **2** could be rationalized to the dissimilar configurations of the stereogenic carbons of the side chains.

**Figure 2.** Selected 2D NMR correlations of **1**

The OH configurations of two stereogenic centers in **1** and **2** were determined by comparison with the ^{13}C NMR data of the structural similar compounds, *erythro*- and *threo*-honokitiol¹⁶ and *erythro*- and *threo*-1-C-syringylglycerol.¹⁷ The difference of ^{13}C NMR chemical shifts of C-7 (δ_{C} 76.0) and C-8 (δ_{C} 76.2) in *erythro*-honokitiol is $\Delta\delta_{\text{C-8-C-7}}$ 0.2, which is smaller than that in *threo*-honokitiol ($\Delta\delta_{\text{C-8(77.8)-C-7(74.6)}}$ 3.2).¹⁶ In addition, the difference of ^{13}C NMR chemical shifts of C-7 (δ_{C} 74.1) and C-8 (δ_{C} 75.3) in *erythro*-1-C-syringylglycerol is $\Delta\delta_{\text{C-8-C-7}}$ 1.2, which is also smaller than that in its *threo*-isomer ($\Delta\delta_{\text{C-8(75.8)-C-7(72.9)}}$ 2.9).¹⁷ Therefore, the relative configurations of **1** and **2** could also be determined base on the rules as mentioned in the above cases.^{16,17} In compound **2**, the difference of the carbon signals of C-7 (δ_{C} 75.8) and C-8 (δ_{C} 76.5) is $\Delta\delta_{\text{C-8-C-7}}$

0.7, which is smaller than that in **1** ($\Delta\delta_{\text{C-8(76.9)-C-7(75.4)}}$ 1.5). Thus, the structures of **1** and **2** were determined as *threo*- and *erythro*- configurations, respectively. As for the absolute configuration of C-7 and C-8, there should be four possibilities, i.e. (7*R*, 8*R*), (7*R*, 8*S*), (7*S*, 8*R*), and (7*S*, 8*S*). In the literature, four stereoisomers of 1-phenylglycidol with (1*R*, 2*R*), (1*R*, 2*S*), (1*S*, 2*R*), (1*S*, 2*S*) configurations were synthesized by an asymmetric method, and the optical data were $[\alpha]_{\text{D}}^{18} - 45.5$ (*c* 3.3), $[\alpha]_{\text{D}}^{25} - 38.5$ (*c* 2.2), $[\alpha]_{\text{D}}^{20} + 37.2$ (*c* 3.4), $[\alpha]_{\text{D}}^{21} + 45.0$ (*c* 2.4), respectively.¹⁸ By comparison of their optical data with those of *threo* isomer (**1**, $[\alpha]_{\text{D}}^{23.5} - 11.0$) and *erythro* isomer (**2**, $[\alpha]_{\text{D}}^{20} - 14.6$), it was confirmed that the configuration of C-7 and C-8 in **1** and **2** were (7*R*, 8*R*) and (7*R*, 8*S*), respectively.

Some chemical constituents from the Magnoliaceae family were reported to have obvious cytotoxic activities.¹⁹ Therefore, all compounds except compound **2** were assayed for their cytotoxicity against the HL-60, SMMC-7721, A-549, MCF-7, and SW-480 human tumor cell lines by the MTT method with *cis*-platin as positive control.¹⁹ Compound **3** showed weak cytotoxic activity against the HL-60 human tumor cell lines with the IC_{50} values of 23.5 μM . The other compounds did not show cytotoxic activity ($\text{IC}_{50} > 40 \mu\text{M}$).

Experimental Section

General Experimental Procedures. Optical rotations were measured with a JASCO DIP-370 digital polarimeter. UV spectra were obtained using a Shimadzu UV-2401A spectrophotometer. A BioRad FtS-135 spectrophotometer was used for scanning IR spectroscopy with KBr pellets. 1D and 2D NMR spectra were recorded on Bruker AM-400, DRX-500 and Bruker Avance III-600MHz spectrometers. Unless otherwise specified, chemical shifts (δ) were expressed in ppm with reference to the solvent signals. High resolution electrospray ionization (HRESIMS) were performed on a VG Autospec-3000 spectrometer under 70 eV. Column chromatography was performed using silica gel (200–300 mesh, Qingdao Marine Chemical, Inc., Qingdao, China). Semi-preparative HPLC was performed on an Agilent 1100 liquid chromatograph with a Zorbax SB-C₁₈, 9.4 mm \times 25 cm, column. Preparative HPLC was performed on a Shimadzu LC-8A preparative liquid chromatograph with a Shimadzu PRC-ODS (K) column.

Fractions were monitored by TLC and spots were visualized by heating the silica gel plates sprayed with 10% H₂SO₄ in EtOH.

Plant Material. The leaves and stems of *M. insignis* (Wall.) Bl. was collected in Kunming Botanic Garden, Yunnan Province, China, in August 2007. The specimen was identified by Prof. Xun GONG and a voucher specimen (No. KIB 2007-08-11) has been deposited at the State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences.

Extraction and Isolation. The air-dried plant material of *M. insignis* (6.5 kg) was ground and exhaustively percolated three times with 70% aqueous Me₂CO at room temperature. The solvent was evaporated in *vacuo*, and the combined crude extract was dissolved in H₂O and partitioned with EtOAc. The EtOAc portion (156 g) was subsequently chromatographed on a silica gel column eluting with CHCl₃-Me₂CO (1:0, 9:1, 8:2, 2:1, 1:1, and 0:1) and was combined into six groups (A–F) according to TCL analysis. Fraction B (21g) was chromatographed on MPLC eluting with MeOH/H₂O (30:70–100:0 gradient system), to give fraction B1–B5. And then fraction B3 was further applied to silica gel CC, and eluted with petroleum ether-acetone (20:1, 12:1, 8:1 and 3:1) to give five fractions B3-1–B3-5. Compounds **1** (7.0 mg, *t_R* = 29 min) and **2** (2.1 mg, *t_R* = 32 min) were obtained from fraction B3-2 by semi-preparative HPLC (42% aq. acetonitrile, 3 mL/min). Fraction B3-3 was chromatographed on Sephadex LH-20 (MeOH) and then successively chromatographed on silica gel CC (CHCl₃-ethyl acetate, 25:1, 16:1, 12:1 and 6:1) to give compound **6** (12 mg). Fraction B4 was further purified on silica gel CC (CHCl₃-ethyl acetate, 24:1–8:1) and then chromatographed on Sephadex LH-20 (MeOH) to give six fractions B4-1–B4-6. Fraction B4-2 were further subjected to semi-preparative HPLC (53% aq. MeOH, 3 mL/min) to afford **3** (9.0 mg, *t_R* = 18 min) and **4** (14.6 mg, *t_R* = 24 min). Compound **5** (5.6 mg, *t_R* = 18 min) were purified by semi-preparative HPLC (55% aq. MeOH, 3 mL/min) from fraction B4-3.

Manneoinsigin A (1): yellow gum; [α]_D^{23.5} – 11.0 (*c* 0.23, MeOH); UV (MeOH) λ_{\max} (log ϵ) 289 (3.27), 214 (4.01), 191 (3.76) nm; IR (KBr) ν_{\max} 3424, 3081, 2924, 2853, 1639, 1619, 1496, 1423, 1230, 915, 826 cm^{–1}; ¹H and ¹³C NMR data, see Table 1; negative ESIMS *m/z* 333 [M – H][–] and isotopic peak *m/z* 335 [M – H][–]; negative HRESIMS *m/z* 333.0897 [M – H][–] (calcd for C₁₈H₁₈O₄Cl, 333.0893) and isotopic peak *m/z* 335.0908 [M – H][–] (calcd for C₁₈H₁₈O₄Cl, 335.0864).

Manneoinsigin B (2): yellow gum; [α]_D^{23.5} – 14.6 (*c* 0.11, MeOH); UV (MeOH) λ_{\max} (log ϵ) 289 (3.49), 212 (4.20), 196 (3.84) nm; IR (KBr) ν_{\max} 3426, 3080, 2927, 2856, 1638, 1624, 1496, 1230, 917, 830 cm^{–1}; ¹H and ¹³C NMR data, see Table 1; negative ESIMS *m/z* 333 [M – H][–] and isotopic peak *m/z* 335 [M – H][–]; HREIMS *m/z* 334.0964 [M]⁺ (calcd for C₁₈H₁₉O₄Cl, 334.0972) and isotopic peak *m/z* 336.0961 [M]⁺ (calcd for C₁₈H₁₉O₄Cl, 336.0942).

Cytotoxicity Assay. The following human tumor cell lines were used: HL-60, MMC-7721, A549, MCF-7, and SW480. All cells were cultured in RPMI-1640 or DMEM medium (Hyclone, Logan, UT), supplemented with 10% fetal bovine serum (Hyclone) at 37 °C in a humidified atmosphere with 5% CO₂. Cell viability was assessed by conducting colorimetric measurements of the amount of insoluble formazan formed in living cells based on the reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma, St. Louis, MO).¹⁹ Briefly, 100 μ L of adherent cells was seeded into each well of a 96-well cell culture plate and allowed to adhere for 12 h before drug addition, while suspended cells were seeded just before drug addition, both with an initial density of 1×10^5 cells/mL in 100 μ L of medium. Each cell line was exposed to the test compound at various concentrations in triplicate for 48 h, with cisplatin and paclitaxel (Sigma) as positive controls. After the incubation, MTT (100 μ g) was added to each well, and the incubation continued for 4 h at 37 °C. The cells were lysed with 100 μ L of 20% SDS-50% DMF after removal of 100 μ L of medium. The optical density of the lysate was measured at 595 nm in a 96-well microtiter plate reader (Bio-Rad 680). The IC₅₀ value of each compound was calculated by Reed and Muench's method.²⁰

Electronic Supplementary Material

Supplementary material is available in the online version of this article at <http://dx.doi.org/10.1007/s13659-012-0063-7> and is accessible for authorized users.

Acknowledgments

This project was supported financially by the NSFC (No. 20802082 and 30830115), the projects from the Chinese Academy of Sciences (KSCX2-EW-Q-10 and KSCX1-YW-R-24), the Major State Basic Research Development Program of China (No. 2009CB522300 and 2009CB940900), and the Natural Science Foundation of Yunnan Province (2012FB178 and 20080A007) and the project of recruited top talent of sciences and technology of Yunnan Province (2009C1120).

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] Youn, U. J.; Chen, Q. C.; Jin, W. Y.; Lee, I. S.; Kim, H. J.; Lee, J. P.; Chang, M. J.; Min, B. S.; Bae, K. H. *J. Nat. Prod.* **2007**, *70*, 1687–1689.
- [2] Watanabe, K.; Watanabe, H. Y.; Goto, Y.; Yamamoto, N.; Yoshizaki, M. *Jpn. J. Pharmacol.* **1975**, *25*, 605–607.
- [3] Watanabe, K.; Watanabe, H.; Goto, Y.; Yamaguchi, M.; Yamamoto, N.; Hagino, K. *Planta Med.* **1983**, *49*, 103–108.
- [4] Watanabe, K. *Gendai Toyo Igaku* **1986**, *7*, 54–59.
- [5] Teng, C. M.; Yu, S. M.; Chen, C. C.; Huang, Y. L.; Huang, T. F. *Life Sci.* **1990**, *47*, 1153–1161.
- [6] Bae, E. A.; Han, M. J.; Kim, D. H. *Planta Med.* **1999**, *65*, 442–443.
- [7] Namba, T.; Tsunazuka, M.; Hattori, M. *Planta Med.* **1982**, *44*, 100–106.
- [8] Sohn, E. J.; Kim, C. S.; Kim, Y. S.; Jung, D. H.; Jang, D. S.; Lee,

- Y. M.; Kim, J. S. *Life Sci.* **2007**, *80*, 468–475.
- [9] Fukuyama, Y.; Otsoshi, Y.; Miyoshi, K.; Nakamura, K.; Kodama, M.; Nagasawa, M.; Hasegawa, T.; Okazaki, H.; Sugawara, M. *Tetrahedron* **1992**, *48*, 377–392.
- [10] College, J. N. M. Ed. Chinese Drug Dictionary. *Shanghai Science and Technology Publishing Co.* **1977**, 1628–1630.
- [11] Vardamides, J. C.; Azebaze, A. G. B.; Nkengfack, A. E.; Van Heerden, F. R.; Fomum, Z. T.; Ngando, T. M.; Conrad, J.; Vogler, B.; Kraus, W. *Phytochemistry* **2003**, *62*, 647–650.
- [12] Sugahara, T.; Yamauchi, S.; Kondo, A.; Ohno, F.; Tominaga, S.; Nakashima, Y.; Kishida, T.; Akiyama, K.; Maruyama, M. *Biosci. Biotechnol. Biochem.* **2007**, *71*, 2962–2968.
- [13] Masuda, T.; Akiyama, J.; Fujimoto, A.; Yamauchi, S.; Maekawa, T.; Sone, Y. *Food Chem.* **2010**, *123*, 442–450.
- [14] Luo, X.; Wu, S.; Ma, Y.; Wu, D. *Acta. Bot. Yunnanica* **2001**, *23*, 368–372.
- [15] Deng, X. M.; Cheng, Y. X.; Zhou, J.; Tan, N. H.; Ding, Z. T. *Acta Bot. Yunnanica* **2001**, *23*, 121–125.
- [16] Shen, C. C.; Ni, C. L.; Shen, Y. C.; Huang, Y. L.; Kuo, C. H.; Wu, T. S.; Chen, C. C. *J. Nat. Prod.* **2009**, *72*, 168–171.
- [17] Otsuka, H.; Takeuchi, M.; Inoshiri, S.; Sato, T.; Yamasaki, K. *Phytochemistry* **1989**, *28*, 883–886.
- [18] Takeshita, M.; Yaguchi, R.; Akutsu, N. *Tetrahedron: Asymmetry* **1992**, *3*, 1369–1372.
- [19] Ping, Z. R.; Hongjie, Z.; Zhongwen, L.; Yulin, Z.; Handong, S. *Phytochemistry* **1992**, *31*, 4237–4240.
- [20] Reed, L. J.; Muench, H., In *Am. J. Hyg.* **1938**, *27*, 493–497.